

Using Michigan's IIS to Monitor Zoster Administration Patterns

Lynsey Kimmins, MPH, Cristi Bramer, MPH, Tina Scott
Michigan Department of Health and Human Services
Division of Immunization

AIRA National Meeting, August 13, 2019

Presentation Overview

- Michigan Care Improvement Registry (MCIR) Background
- Timeline of Zoster Vaccines and Recommendations
- Zoster Vaccine Administration Patterns
 - Zoster Dose Reporting
 - Recipient Age at Administration
 - Shingrix Series
- Key Takeaways

MCIR Background

- Started in 1998, lifespan registry since 2006
- Vaccination record submission for adults 20 years and older is strongly encouraged but is not required
 - Adult vaccination data reporting has been increasing in quality and quantity, most notably since the H1N1 pandemic of 2009 and the implementation of HL7 messaging in 2012
 - Most frequent reporters of adult dose to the MCIR, by MCIR facility type, are family practices, pharmacies and internal medicine practices
- As of June 22, 2019, there are **2,408,015 child** (≤ 18) and **7,764,613 adult** (>18) **Michigan** residents in the MCIR

Timeline: Zoster Vaccines

2006

- Zostavax licensed by FDA for adults 60+

2008

- ACIP recommended Zostavax for adults 60+


2011

- FDA licensed Zostavax for adults 50+

**2011 &
2013**

- ACIP considered (but did not recommend) Zostavax for adults 50+

Estimated proportion of adults 60+ who ever received a herpes zoster vaccination— National Health Interview Survey (NHIS), US and MCIR, Michigan, 2013-2017



*Denominator based on Census estimates; CYQ4 estimates for 1+ Zostavax vaccine.

Adult data reporting to the MCIR is strongly encouraged but is not required.

Timeline: Zoster Vaccines

**October
2017**

- Shingrix licensed by FDA for adults 50+
- Rare ACIP preferential recommendation for Shingrix

**January
2018**

- CDC released new zoster recommendations

**April
2018**

- Manufacturer reported Shingrix shortages

Timeline: Zoster Vaccines- Shingrix Shortage

July
2018

- Manufacturer implemented order limits for providers, distributors, and wholesalers that order from the manufacturer directly


August
2018

- CDC released clinical guidance
- *Shingrix is the preferred shingles vaccine*

September
2018


- *Give first consideration to patients due for their second dose of Shingrix*

Zoster doses administered and reported for adults 50+ by vaccine type, month and year, *MCIR, Michigan, January 2013 - May 2019*




*Includes first recorded dose of Zostavax and/or Shingrix recorded in the MCIR

Zoster dose coadministrations reported by select vaccine type, *MCIR, Michigan, May 2006 - May 2019*




Distribution of age at administration for Zostavax doses by year, MCIR, Michigan, 2013-2019




*Age at administration of first dose of Zostavax recorded in the MCIR

Distribution of age at administration for Shingrix doses by month, *MCIR, Michigan, January 2018- May 2019*




*Age at administration of first dose of Shingrix recorded in the MCIR

Zoster doses administered and reported for adults 50+ by facility type and year, *MCIR, Michigan, 2013 – 2019 (January-May)*




*Includes first recorded dose of Zostavax and/or Shingrix recorded in the MCIR

Shingrix doses administered and reported by facility type and week, MCIR, Michigan, January 2018 - May 2019




Shingrix doses administered and reported by facility type and day, Michigan, MCIR, April 1, 2019 - May 31, 2019




Shingrix Series Completion, *Adults ≥50 years, MCIR, Michigan, October 20, 2017 – May 31, 2019*

- Over **195,000 adults** 50+ years have had at least one Shingrix dose reported to the MCIR
 - 71% have had at least one dose reported by a pharmacy location
- According to the MCIR, **57%** of those who have initiated the series have completed the 2-dose series
 - 69% of those individuals completed the series entirely at pharmacy locations
- **57%** of Shingrix recipients were female
- **38%** of Shingrix recipients also have a Zostavax dose in the MCIR


Shingrix doses administered and reported for adults 50+ by first and second dose and month, MCIR, Michigan, January 2015 - May 2019




Intervals between Shingrix doses, *MCIR*, *Michigan, October 20, 2017 – June 8, 2019*




Distribution of intervals between Shingrix doses by month, *MCIR, Michigan, January – December 2018*



Distribution of distance traveled for first Shingrix dose by month, *MCIR, Michigan, March 1, 2018 – May 31, 2019*



Distribution of distance traveled for first Shingrix dose by facility type, *MCIR, Michigan, January 1, 2018 – May 31, 2019*



Key Takeaways

- *Pharmacies* continue to be the top reporters of Zoster vaccines
- Shingrix dose reporting patterns have been *variable* since its introduction
- Shingrix age at administration, interval between doses and distance traveled have remained fairly *stable* over the past year and a half
- *Over half* of Michigan residents that have initiated the Shingrix series have completed the series in the recommended timeframe
- *Established* adult reporting to the MCIR has allowed for better tracking of Shingrix uptake and administration patterns